

MEASUREMENT OF $\sin^2 \theta_W$ IN νN SCATTERING FROM NuTeV

R. A. JOHNSON¹, T. ADAMS⁴, A. ALTON⁴, S. AVVAKUMOV⁷,
 L. DE BARBARO⁵, P. DE BARBARO⁷, R. H. BERNSTEIN³, A. BODEK⁷,
 T. BOLTON⁴, J. BRAU⁶, D. BUCHHOLZ⁵, H. BUDD⁷, L. BUGEL³,
 J. CONRAD², R. B. DRUCKER⁶, R. FREY⁶, J. GOLDMAN⁴,
 M. GONCHAROV⁴, D. A. HARRIS⁷, S. KOUTSOLIOTAS^{2,*}, J. H. KIM^{2, α} ,
 M. J. LAMM³, W. MARSH³, D. MASON⁶, C. MCNULTY², K. S. MCFARLAND⁷,
 D. NAPLES^{4,+}, P. NIENABER³, A. ROMOSAN², W. K. SAKUMOTO⁷,
 H. SCHELLMAN⁵, M. H. SHAEVITZ², P. SPENTZOURIS^{2,**}, E. G. STERN^{2, $\dagger$} ,
 M. VAKILI^{1, $\parallel$} , A. VAITAITIS², V. WU¹, U. K. YANG⁷, J. YU³, and
 G. P. ZELLER⁵

¹University of Cincinnati, Cincinnati, OH 45221, ²Columbia University, New York, NY 10027, ³Fermi National Accelerator Laboratory, Batavia, IL 60510, ⁴Kansas State University, Manhattan, KS 66506, ⁵Northwestern University, Evanston, IL 60208, ⁶University of Oregon, Eugene, OR 97403, ⁷University of Rochester, Rochester, NY 14627

Current Addresses: *Bucknell University, ^{$\dagger$} Lawrence Berkeley National Laboratory, ^{$\parallel$} Texas A&M University, **Fermi National Accelerator Laboratory, +University of Pittsburgh, ^{α} University of California at Irvine, ^{$\dagger$} Lucent Technologies

Presented by Randy Johnson

We report the measurement of $\sin^2 \theta_w$ in νN deep inelastic scattering from the NuTeV experiment. By using separate neutrino and anti-neutrino beams, NuTeV is able to determine $\sin^2 \theta_W$ with smaller systematic and similar or smaller statistical errors when compared to previous neutrino experiments. NuTeV measures $\sin^2 \theta_W^{(\text{on-shell})} = 0.2253 \pm 0.0019(\text{stat.}) \pm 0.0010(\text{sys.})$, which implies $M_W = 80.26 \pm 0.11 \text{ GeV}/c^2$.

1 Introduction

Neutrino experiments play a key role in establishing the validity of the electroweak Standard Model. Even today with the large samples of W and Z boson events, precision νN experiments are still competitive in determining the electroweak model parameters and also show that the model is valid over many orders of magnitude in q^2 . To date, neutrino experiments have used the Llewellyn Smith¹ relationship to determine $\sin^2 \theta_W$.^{2,3} However, the dominate systematic error, the experimental determination of the effective charm mass, has limited the precision of these measurements. Paschos and Wolfenstein⁴

have shown that if the neutral and charged current cross sections for neutrinos and anti-neutrinos could be measured separately, then the ratio

$$\begin{aligned} R^- &= \frac{\sigma(\nu_\mu N \rightarrow \nu_\mu X) - \sigma(\bar{\nu}_\mu N \rightarrow \bar{\nu} X)}{\sigma(\nu_\mu N \rightarrow \mu^- X) - \sigma(\bar{\nu}_\mu N \rightarrow \mu^+ X)} \\ &= \frac{R^\nu - r R^{\bar{\nu}}}{1 - r} = \frac{1}{2} - \sin^2 \theta_W = g_L^2 - g_R^2 \end{aligned} \quad (1)$$

(where $R^\nu = \frac{\sigma(\nu_\mu N \rightarrow \nu_\mu X)}{\sigma(\nu_\mu N \rightarrow \mu^- X)}$, $R^{\bar{\nu}} = \frac{\sigma(\bar{\nu}_\mu N \rightarrow \bar{\nu} X)}{\sigma(\bar{\nu}_\mu N \rightarrow \mu^+ X)}$, $r = \frac{\sigma(\bar{\nu}_\mu N \rightarrow \mu^+ X)}{\sigma(\nu_\mu N \rightarrow \mu^- X)}$, $g_{L,R}^2 = u_{L,R}^2 + d_{L,R}^2$ = the sum of the squares of the quark couplings, and the relationship to $\sin^2 \theta_W$ is a tree level relationship only) eliminates the effect of the sea quarks on the determination of $\sin^2 \theta_W$ and, hence, the majority of the error from the mass of the charm quark. To make such a measurement was a goal of the NuTeV experiment at Fermilab.

2 NuTeV experiment

In order to measure R^- , the cross sections for νN and $\bar{\nu} N$ scattering must be measured separately. For NuTeV, we designed and built a Sign Selected Quadrupole Train (SSQT) which selected the charge of the mesons that were directed into our decay volume, and, thus, selected whether neutrinos or anti-neutrinos hit the detector. The fluxes of neutrino varieties in neutrino and anti-neutrino modes are shown in Fig. 1(a). The wrong sign contamination fraction was less than 1×10^{-3} in neutrino mode and 2×10^{-3} in anti-neutrino mode. The electron neutrino contamination in the beam is a serious background in the determination of R^- . The majority of ν_e 's come from charged kaon decays which can be kinematically related to the high energy ν_μ spectrum that comes from the same particles. The design of the SSQT greatly reduced the contribution of the neutral kaon decays to the ν_e flux in comparison to other experiments.

The NuTeV experiment used the refurbished CCFR iron calorimeter and toroid spectrometer that is described elsewhere.⁵ Data were collected during the '96-'97 Fermilab fixed target run. Charged current ν and $\bar{\nu}$ events were differentiated from neutral current events by the length in the calorimeter over which energy was deposited. Long events indicate the presence of a muon and are classified as charged current events; neutral current events only contain a hadronic shower and are therefore short. The length distribution for neutrino and anti-neutrino events is shown in Fig. 1(b). A detailed Monte Carlo program was used to determine the contamination of neutral current events in the long distribution and of charged current events in the short distribution. The




Figure 1: (a) *Preliminary* NuTeV Neutrino and Anti-Neutrino Fluxes; (b) *Preliminary* NuTeV Event Length Distribution.

detector parameters used in this program were measured using a test beam which constantly monitored and calibrated the detector.

3 Results

Rather than measuring R^- directly, we chose to measure a “pseudo” R^-

$$\text{pseudo}R^- = R^\nu - \alpha R^{\bar{\nu}} \quad (2)$$

where $\alpha = 0.5136$ was chosen to minimize the effects of the charmed quark mass on our result given the differences in fluxes and cross sections between the ν and $\bar{\nu}$ modes. We varied the value of $\sin^2 \theta_W$ in our Monte Carlo until the predicted value of $\text{pseudo}R^-$ matched the measured one.

The preliminary result from the NuTeV data is

$$\begin{aligned} \sin^2 \theta_W^{(\text{on-shell})} &= 0.2253 \pm 0.0019(\text{stat.}) \pm 0.0010(\text{sys.}) \\ &- 0.00142 \times \left(\frac{M_{\text{top}}^2 - (175 \text{ GeV}/c^2)^2}{(100 \text{ GeV}/c^2)^2} \right) + 0.00048 \times \log_e \left(\frac{M_{\text{Higgs}}^2}{150 \text{ GeV}/c^2} \right). \end{aligned} \quad (3)$$

The small residual dependence of our result on M_{top} and M_{Higgs} comes from the leading terms in the electroweak radiative corrections⁶. The sources of the statistical and systematic errors for this result are given in Table 1.

Table 1: Sources of uncertainties in the determination of $\sin^2 \theta_W$.

SOURCE OF UNCERTAINTY		$\delta \sin^2 \theta_W$
<i>Statistics:</i>	Data	0.00188
	Monte Carlo	0.00028
TOTAL STATISTICS		0.00190
<i>$\nu_e/\bar{\nu}_e$ Flux:</i>	$K^\pm$ (1.1%)	0.00024
	Other sources of ν_e 's	0.00048
<i>Energy Measurement:</i>	Calibrations (0.5%)	0.00043
	Muon Energy Deposition (3%)	0.00004
	Energy Resolution	0.00004
<i>Event Length:</i>	Hadron Shower	0.00015
	Longitudinal Vertex Determination	0.00015
	Counter Edge Location	0.00010
	Counter Efficiency and Noise	0.00016
TOTAL EXP. SYST.		0.00075
<i>Sea Quarks:</i>	Strange Sea	0.00034
	V_{cd}	0.00004
	Charm Sea	0.00009
	Charm Mass	0.00009
<i>Other $\nu/\bar{\nu}$ Cross-Section Differences: $\sigma^\nu/\sigma^{\bar{\nu}}$</i>		0.00023
	Non-Isoscalar Target	0.00013
	Radiative Corrections	0.00051
<i>Non-QPM Cross-Section:</i>	Higher Twist	0.00013
	Longitudinal Structure Function	0.00007
TOTAL PHYSICS MODEL		0.00070
TOTAL UNCERTAINTY		0.0022

Since $\sin^2 \theta_W^{(\text{on-shell})} = 1 - M_W^2/M_Z^2$, the result given in Eq. 3 is equivalent to

$$M_W = 80.26 \pm 0.10(\text{stat.}) \pm 0.05(\text{sys.}) \\ + 0.073 \times \left(\frac{M_{\text{top}}^2 - (175 \text{ GeV}/c^2)^2}{(100 \text{ GeV}/c^2)^2} \right) - 0.025 \times \log_e \left(\frac{M_{\text{Higgs}}^2}{150 \text{ GeV}/c^2} \right). \quad (4)$$

This result is compared with other measurements of the W mass in Fig. 2(a).

It is possible to extract the NuTeV result in a model independent framework in terms of the left and right handed quark couplings. For our data

$$0.4530 - \sin^2 \theta_W = 0.2277 \pm 0.0022 \\ = 0.8587u_L^2 + 0.8828d_L^2 - 1.1657u_R^2 - 1.2288d_R^2. \quad (5)$$

(Note the similarity of this with Eq. 1.) This result is plotted on the g_L^2 - g_R^2 plane in Fig. 2(b) and compared with the CCFR result which measured




Figure 2: (a) Comparison of the measured M_W mass from NuTeV and other experiments; (b) Comparison of the model independent couplings between NuTeV and CCFR (Point is the '98 Standard Model fit of world data by the Particle Data Group⁷).

$R^\nu = g_L^2 + 0.64g_R^2$. The results of the two experiments are consistent with the Standard Model.

Acknowledgments

We gratefully acknowledge the substantial contributions in the construction and operation of the NuTeV beamlines and the refurbishment of the NuTeV detector made by the staff of the Fermilab Beams and Particle Physics Division. This work was supported by grants from the U.S. Department of Energy, from the National Science Foundation, and from the Alfred P. Sloan Foundation.

References

1. C.H. Llewellyn Smith, *Nucl. Phys. B* **228**, 205 (1983).
2. K.S. McFarland, *et al.*, *Eur. Phys. Journal* **C1**, 509 (1998).
3. A. Blondel, *et al.*, *Z. Phys. C* **45**, 361 (1990); J. Allaby, *et al.*, *Z. Phys. C* **36**, 611 (1985).
4. E.A. Paschos and L. Wolfenstein, *Phys. Rev. D* **7**, 91 (1973).
5. W.S. Sakumoto, *et al.*, *Nucl. Instrum. Methods A* **294**, 179 (1991); B.J. King, *et al.*, *Nucl. Instrum. Methods A* **302**, 254 (1991).
6. D.Yu. Bardin and V.A. Dokuchaeva, JINR-E2-86-2, 60(1986).
7. Particle Data Group: C. Caso, *et al.*, *Eur. Phys. Journal* **C3**, 97 (1998).